

Citation for published version:

Bill, A, Guiver, C, Logemann, H & Townley, S 2014, A stability/instability trichotomy for non-negative Lur'e systems. in *Proceedings of the MTNS 2014*. <<http://fwn06.housing.rug.nl/mtns2014-papers/extendedAbstracts/0247.pdf>>

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Publisher Rights
CC BY

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

A stability/instability trichotomy for non-negative Lur'e systems*

Adam Bill¹, Chris Guiver², Hartmut Logemann¹ and Stuart Townley²

Abstract—We identify a stability/instability trichotomy for a class of non-negative continuous-time Lur'e systems. Asymptotic as well as input-to-state stability concepts (ISS) are considered. The presented trichotomy rests on Perron-Frobenius theory, absolute stability theory and recent ISS results for Lur'e systems.

I. INTRODUCTION

Let $A = (a_{ij}) \in \mathbb{R}^{n \times n}$ and $b, c \in \mathbb{R}^n$ and consider the corresponding single-input single-output non-negative linear system

$$\dot{x} = Ax + bu, \quad x(0) = \xi \in \mathbb{R}_+^n; \quad y = c^T x. \quad (1)$$

We assume that

(A1) A is Metzler, $b, c \in \mathbb{R}_+^n$ and $b, c \neq 0$ holds.

We recall that $A = (a_{ij})$ is *Metzler* if $a_{ij} \geq 0$ for $i \neq j$ (all off-diagonal elements are non-negative).

System (1) is said to be *non-negative* if (A1) holds and $u \geq 0$. Non-negative systems of the form (1) occur naturally in biological, ecological and economic contexts.

We impose the following assumptions.

(A2) A is Hurwitz.

(A3) There exist non-negative numbers α and κ such that $\alpha I + A + \kappa bc^T$ is primitive.

Recall that (A3) means that the matrix $(\alpha I + A + \kappa bc^T)^k$ is a positive matrix for some $k \in \mathbb{N}$.

In the following, let G denote the transfer function of (1), that is, $G(s) := c^T(sI - A)^{-1}b$.

Lemma 1.1: Assume that (A1)-(A3) hold. Then $G(0) > 0$ and $\|G\|_{H^\infty} = G(0)$.

A proof of Lemma 1.1 can be found in [1].

Applying nonlinear non-negative feedback $u = f(y)$ to (1), where $f: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is locally Lipschitz, leads to the following non-negative Lur'e system

$$\dot{x} = Ax + bf(c^T x), \quad x(0) = \xi \in \mathbb{R}_+^n. \quad (2)$$

We assume that the following assumption holds.

(A4) $f: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is locally Lipschitz and $f(0) = 0$.

Whilst absolute stability of Lur'e systems is a classical topic in control theory (see, for example, [2], [3], [8]), it seems that non-negative Lur'e systems have not received

much attention (see however [7] which provides an analysis of the stability properties of a class of non-negative discrete-time Lur'e systems).

Assuming that (A1)-(A4) hold, we set

$$p := \frac{1}{G(0)},$$

and consider the following three cases.

Case 1. $f(z)/z \leq p$ for all $z > 0$.

Case 2. $\inf_{z>0} f(z)/z > p$.

Case 3. There exists $y^* > 0$ such that $f(y^*) = py^*$ and

$$\left| \frac{f(z) - f(y^*)}{z - y^*} \right| \leq p \text{ for all } z > 0, z \neq y^*.$$

The condition in Case 3 means that the graph of f is “sandwiched” between the straight lines l_1 and l_2 given by $l_1(z) = pz$ and $l_2(z) = 2py^* - pz$, see Figure 1.




Fig. 1. Case 3: graph of f “sandwiched” between the lines l_1 and l_2 .

II. LYAPUNOV STABILITY RESULTS

In this section, we present results which describe the stability/instability properties in each of three cases, where “stability” is interpreted in the sense of Lyapunov.

Let $x(\cdot; \xi)$ denote the unique maximally defined forward solution of (2) with maximal interval of existence $[0, \omega_\xi)$, where $0 < \omega_\xi \leq \infty$.

The proposition below relates to Case 1. It follows from well known results in absolute stability theory, see, for example, [3].

Proposition 2.1: Assume that (A1)-(A4) hold.

(a) If $f(z)/z \leq p$ for all $z > 0$, then the equilibrium 0 is stable in the large in the sense that there exists $\Gamma \geq 1$ such that, for every $\xi \in \mathbb{R}_+^n$, $\omega_\xi = \infty$ and

$$\|x(t; \xi)\| \leq \Gamma \|\xi\| \quad \forall t \geq 0.$$

*This work was supported by EPSRC (Grant EP/I019456/1)

¹ Department of Mathematical Sciences, University of Bath, Bath BA2 7AY, UK A.R.Bill@bath.ac.uk, H.Logemann@bath.ac.uk

² Environment and Sustainability Institute, University of Exeter, Penryn Campus, Penryn, Cornwall, TR10 9EZ, UK C.Guiver@exeter.ac.uk, S.B.Townley@ex.ac.uk

(b) If $f(z)/z < p$ for all $z > 0$, then the equilibrium 0 is globally asymptotically stable. In particular, for every $\xi \in \mathbb{R}_+^n$, $\omega_\xi = \infty$ and $x(t; \xi) \rightarrow 0$ as $t \rightarrow \infty$.

(c) If $\sup_{z>0} f(z)/z < p$, then the equilibrium 0 is globally exponentially stable, that is, there exist $N \geq 1$ and $\nu > 0$ such that, for every $\xi \in \mathbb{R}_+^n$, $\omega_\xi = \infty$ and

$$\|x(t; \xi)\| \leq Ne^{-\nu t} \|\xi\| \quad \forall t \geq 0.$$

In Case 2, the solutions of (2) diverge to ∞ for every non-zero initial condition. More precisely, we have the following result.

Theorem 2.2: Assume that (A1)-(A4) hold and $\inf_{z>0} f(z)/z > p$. Let $\xi \in \mathbb{R}_+^n$, $\xi \neq 0$, be such that the solution $x(t; \xi)$ exists for all $t \geq 0$. Then

$$\lim_{t \rightarrow \omega_\xi} x_i(t; \xi) = \infty \quad \forall i \in \{1, \dots, n\},$$

where $x_i(\cdot; \xi)$ denotes the i -th component of $x(\cdot; \xi)$.

We proceed to consider Case 3.

Theorem 2.3: Assume that (A1)-(A4) hold.

(a) If there exists $y^* > 0$ such that $f(y^*) = py^*$ and

$$\left| \frac{f(z) - f(y^*)}{z - y^*} \right| \leq p \quad \forall z \geq 0, z \neq y^*$$

then $x^* = -pA^{-1}by^* \in \mathbb{R}_+^n$ is an equilibrium of (2) and x^* is stable in the large in the sense that there exists $\Gamma \geq 1$ such that, for every $\xi \in \mathbb{R}_+^n$, $\omega_\xi = \infty$ and

$$\|x(t; \xi) - x^*\| \leq \Gamma \|\xi - x^*\| \quad \forall t \geq 0.$$

(b) If there exists $y^* > 0$ such that $f(y^*) = py^*$ and

$$\left| \frac{f(z) - f(y^*)}{z - y^*} \right| < p \quad \forall z > 0, z \neq y^*$$

then 0 and $x^* = -pA^{-1}by^* \in \mathbb{R}_+^n$ are the only equilibria of (2) and x^* is globally asymptotically stable in the sense that x^* is stable in the large (see statement (a) of this theorem) and, for every $\xi \in \mathbb{R}_+^n$ such that $\xi \neq 0$, $\omega_\xi = \infty$ and $x(t; \xi) \rightarrow x^*$ as $t \rightarrow \infty$.

(c) If there exists $y^* > 0$ such that $f(y^*) = py^*$,

$$\left| \frac{f(z) - f(y^*)}{z - y^*} \right| < p \quad \forall z > 0, z \neq y^*$$

and

$$\limsup_{y \rightarrow y^*} \left| \frac{f(z) - f(y^*)}{y - y^*} \right| < p,$$

and if

$$\liminf_{z \rightarrow 0} \frac{f(z)}{z} > p, \quad (3)$$

then 0 and $x^* = -pA^{-1}by^* \in \mathbb{R}_+^n$ are the only equilibria of (2) and x^* is “semi-globally” exponentially stable in the sense that, for every compact set $K \subset \mathbb{R}_+^n$ with $0 \notin K$, there exists $N \geq 1$ and $\nu > 0$ such that, for every $\xi \in K$, $\omega_\xi = \infty$ and

$$\|x(t; \xi) - x^*\| \leq Ne^{-\nu t} \|\xi - x^*\| \quad \forall t \geq 0.$$

(d) If (3) holds and there exists $y^* > 0$ such that $f(y^*) = py^*$ and, for every $\varepsilon > 0$,

$$\sup_{z \geq \varepsilon, z \neq y^*} \left| \frac{f(z) - f(y^*)}{z - y^*} \right| < p,$$

then 0 and $x^* = -pA^{-1}by^* \in \mathbb{R}_+^n$ are the only equilibria of (2) and x^* is “quasi-globally” exponentially stable in the sense that, for every $\delta > 0$ there exist $N \geq 1$ and $\nu > 0$ such that, for every $\xi \in \mathbb{R}_+^n$ with $\|\xi\| \geq \delta$, $\omega_\xi = \infty$ and

$$\|x(t; \xi) - x^*\| \leq Ne^{-\nu t} \|\xi - x^*\| \quad \forall t \geq 0. \quad (4)$$

We remark that “global” exponential stability of x^* (in the sense that there exist $N \geq 1$ and $\nu > 0$ such that (4) is satisfied for all $\xi \in \mathbb{R}_+^n$ with $\xi \neq 0$) does not hold. This is an immediate consequence of the following result which follows from continuity properties of the flow generated by the Lur’e system (2).

Proposition 2.4: Assume that (A1)-(A4) hold and that there exists $y^* > 0$ such that $f(y^*) = py^*$. Then, for every sequence (t_n) in \mathbb{R}_+ with $t_n \rightarrow \infty$ as $n \rightarrow \infty$, there exists a sequence (ξ_n) in \mathbb{R}_+^n with $\xi_n \neq 0$ and $\xi_n \rightarrow 0$ as $n \rightarrow \infty$ and such that

$$\lim_{n \rightarrow \infty} \frac{\|x(t_n; \xi_n) - x^*\|}{\|\xi_n - x^*\|} = 1,$$

where $x^* = -pA^{-1}by^*$.

Discrete-time results similar to statement (b) of Proposition 2.1, Theorem 2.2 and statement (b) of Theorem 2.3 can be found in [7].

Proofs of the results in Section II can be found in [1].

III. INPUT-TO-STATE STABILITY RESULTS

Finally, we investigate the stability behaviour of (2) subject to non-negative disturbances, that is, we analyze input-to-state stability (ISS) properties of the forced Lur’e system

$$\dot{x} = Ax + b(f(c^T x) + d), \quad x(0) = \xi \in \mathbb{R}_+^n, \quad (5)$$

where $d : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is locally essentially bounded. The unique maximally defined forward solution of (5) is denoted by $x(\cdot; \xi, d)$.

For an overview of ISS theory, the reader is referred to [6]. We recall some terminology and notation relating to comparison functions. Let \mathcal{K} denote the set of all continuous functions $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that $\varphi(0) = 0$ and φ is strictly increasing. Moreover, define $\mathcal{K}_\infty := \{\varphi \in \mathcal{K} : \lim_{s \rightarrow \infty} \varphi(s) = \infty\}$. We denote by \mathcal{KL} the set of functions in two variables $\psi : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with the following properties: $\psi(\cdot, t) \in \mathcal{K}$ for all $t \geq 0$, and $\psi(s, \cdot)$ is nonincreasing with $\lim_{t \rightarrow \infty} \psi(s, t) = 0$ for all $s \geq 0$.

The following proposition is a consequence of recent ISS results for Lur’e systems, see [4], [5].

Proposition 3.1: Assume that (A1)-(A4) hold. If there exists $\rho \in \mathcal{K}_\infty$ such that

$$f(z) \leq pz - \rho(z) \quad \forall z \geq 0,$$

then the equilibrium 0 of the unforced Lur'e system (2) is ISS in the sense that there exist $\psi \in \mathcal{KL}$ and $\varphi \in \mathcal{K}$ such that for all $\xi \in \mathbb{R}_+^n$ and all non-negative $d \in L_{\text{loc}}^\infty(\mathbb{R}_+)$, $x(\cdot; \xi, d)$ is defined on \mathbb{R}_+ and

$$\|x(t; \xi, d)\| \leq \psi(\|\xi\|, t) + \varphi(\|d\|_{L^\infty(0,t)}) \quad \forall t \geq 0.$$

The following theorem shows that, under suitable assumptions, the equilibrium x^* has stability properties which are similar to ISS.

Theorem 3.2: Assume that (A1)-(A4) hold and that there exists $y^* > 0$ such that $f(y^*) = py^*$ and

$$\left| \frac{f(z) - f(y^*)}{z - y^*} \right| < p \quad \forall z > 0, z \neq y^*, \quad (6)$$

Furthermore, assume that (3) holds and

$$pz - f(z) \rightarrow \infty \quad \text{as } z \rightarrow \infty. \quad (7)$$

Then 0 and $x^* = -pA^{-1}by^* \in \mathbb{R}_+^n$ are the only equilibria of the unforced Lur'e system (2) and x^* is “quasi ISS” in the sense that, for every $\delta > 0$, there exist $\psi \in \mathcal{KL}$ and $\varphi \in \mathcal{K}$ such that for all $\xi \in \mathbb{R}_+^n$ with $\|\xi\| \geq \delta$ and all non-negative $d \in L_{\text{loc}}^\infty(\mathbb{R}_+)$, $x(\cdot; \xi, d)$ is defined on \mathbb{R}_+ and

$$\|x(t; \xi, d) - x^*\| \leq \psi(\|\xi - x^*\|, t) + \varphi(\|d\|_{L^\infty(0,t)}) \quad \forall t \geq 0. \quad (8)$$

To relate the conditions (6) and (7) to those in Proposition 3.1, we note that if (6) and (7) hold, then, for every $\varepsilon > 0$, there exists $\rho \in \mathcal{K}_\infty$ such that

$$|f(z) - f(y^*)| \leq p|z - y^*| - \rho(|z - y^*|) \quad \forall z \geq \varepsilon, z \neq y^*.$$

The proof of Theorem 3.2 is based on Proposition 3.1 and the following lemma.

Lemma 3.3: Assume that (A1)-(A4) hold. If (3) is satisfied and there exists $y^* > 0$ such that $f(y^*) = py^*$ and (6) holds, then, for every $\delta > 0$, there exist constants $\eta > 0$ and $\tau \geq 0$ such that for all $\xi \in \mathbb{R}_+^n$ with $\|\xi\| \geq \delta$ and all non-negative $d \in L_{\text{loc}}^\infty(\mathbb{R}_+)$, $x(\cdot; \xi, d)$ is defined on \mathbb{R}_+ and

$$c^T x(t; \xi, d) \geq \eta \quad \forall t \geq \tau.$$

This lemma also plays a key roll in the proof of statements (b)-(d) of Theorem 2.3 (with disturbance $d = 0$). Detailed proofs of Proposition 3.1, Theorem 3.2 and Lemma 3.3 can be found in [1].

Finally, it follows from Proposition 2.4 that “global” ISS of x^* (in the sense that there exist $\psi \in \mathcal{KL}$ and $\varphi \in \mathcal{K}$ such that (8) is satisfied for all $\xi \in \mathbb{R}_+^n$ with $\xi \neq 0$ and all non-negative $d \in L_{\text{loc}}^\infty(\mathbb{R}_+)$) does not hold.

REFERENCES

- [1] A. Bill, C. Guiver, H. Logemann and S. Townley. Stability of non-negative Lur'e systems, in preparation, 2014.
- [2] W.M. Haddad and V. Chellaboina. *Nonlinear Dynamical Systems and Control*, Princeton University Press, Princeton, 2008.
- [3] B. Jayawardhana, H. Logemann and E.P. Ryan. The circle criterion and input-to-state stability: new perspectives on a classical result, *IEEE Control Systems Magazine*, **31**, 32-67, (August 2011).
- [4] E. Sarkans and H. Logemann. Input-to-state stability for Lur'e systems, in preparation, 2013.
- [5] E. Sarkans. Absolute stability and input-to-state stability of Lur'e systems, PhD thesis, in preparation, University of Bath, 2014.
- [6] E. D. Sontag. Input to state stability: basic concepts and results, in *Nonlinear and Optimal Control Theory*, P. Nistri and G. Stefani, Eds. Berlin: Springer-Verlag, 2006, pp. 163-220.
- [7] S. Townley, R. Rebarber and B. Tenhumberg. Feedback control systems analysis of density dependent population dynamics, *Systems & Control Letters*, **61**, 309-315, 2012.
- [8] M. Vidyasagar. *Nonlinear Systems Analysis*, 2nd ed., Prentice-Hall, Englewood Cliffs, NJ, 1993.